

Variance Decomposition Techniques for Sensitivity Analysis in Stochastic / Noisy Models with Uncertain Parameters

Olivier Le Maître^{1,2,3}, Omar Knio^{1,2}

1- Duke University, Durham, North Carolina

2- KAUST, Saudi-Arabia

3- LIMSI CNRS, Orsay, France



UNCECOMP, Crete

1 Variance-Based Sensitivity Analysis for SODE's

- Variance decomposition
- Galerkin approximation
- Examples

2 Stochastic Simulations

- stochastic simulators
- Variance Decomposition
- Examples

3 Conclusion

Stochastic models

Physical systems with

- complex **small scale** dynamics (MD, chemical systems, ...)
- **random forcing** and source terms (finance, wind-load, ...)
- **unresolved scales** (turbulence, climate modeling, ...)

are often tackled by means of **stochastic modeling** where complex / unknown / unresolved phenomenons are accounted for by the introduction of noisy dynamics.

In addition to the effect of the noise, the model may involve **unknown parameters** : e.g. noise level, physical constants and parameters, initial conditions, ...

Our general objective is to **propagate / assess the impact of parameters uncertainty** within such stochastic models while characterizing the effect of **inherent noise** :

global sensitivity analysis & analysis of the variance

Stochastic ODEs

We consider a simple systems driven by random noise (Ito equation) : for $t \in [0, T] \doteq \mathcal{T}$

$$dX(t) = C(X(t))dt + D(X(t))dW(t), \quad X(t=0) = X_0,$$

where

- $X(t) \in \mathbb{R}$ is the solution,
- $W(t)$ is **the Wiener process**,
- $C(\cdot)$ is the drift function,
- and $D(\cdot)$ is the diffusion coefficient.

The solution can be computed through **MC simulation**, solving (e.g.)

$$X_{i+1} = X_i + C(X_i)\Delta t + D(X_i)\Delta W_i, \quad X_i \approx X(i\Delta t),$$

drawing **iid random variables** $\Delta W_i \sim N(0, \Delta t)$.

Sample estimate expectation, moments, quantiles, probability law, . . . , of the stochastic process $X(t)$:

$$\mathbb{E} \{g(X(i\Delta t))\} \approx \frac{1}{M} \sum_{l=1}^M g(X_l^i).$$

Stochastic ODEs with parametric uncertainty

The drift function and diffusion coefficient can involve some uncertain parameters Q :

$$dX(t) = C(X(t); Q)dt + D(X(t); Q)dW(t), \quad X(t=0) = X_0.$$

We consider that :

- Q random with known probability law,
- Q and W are assumed independent.

The solution can be seen as a functional of $W(t)$ and Q : $X(t) = X(t, W, Q)$. We shall assume, $\forall t \in \mathcal{T}$,

- 1 $\mathbb{E} \{X^2\} < \infty$,
- 2 $\mathbb{E} \{ \mathbb{E} \{X|W\}^2 \} \doteq \mathbb{E} \{X_{|W}^2\} < \infty$,
- 3 $\mathbb{E} \{ \mathbb{E} \{X|Q\}^2 \} \doteq \mathbb{E} \{X_{|Q}^2\} < \infty$.

We want to investigate the respective impact of Q , W on the dynamics.

Classical sensitivity analysis

Focusing on the two first moments, global SA for the random parameters Q is based on :

- 1 approximating the mean and variance of $X|_Q$

$$\mathbb{E}\{X|_Q\} = \mu_X(Q), \quad \mathbb{V}\{X|_Q\} = \Sigma_X^2(Q),$$

- 2 perform a GSA of $\mu_X(Q)$ and $\Sigma_X^2(Q)$ with respect to the input parameters in Q .

In particular, for independent parameters Q , Polynomial Chaos approximations :

$$\mu_X(Q) \approx \sum_{\alpha} \mu_{\alpha} \psi_{\alpha}(Q), \quad \Sigma_X^2(Q) \approx \sum_{\alpha} \Sigma_{\alpha}^2 \psi_{\alpha}(Q).$$

PC expansion coefficients can be computed / estimated by means of **Non-Intrusive Spectral Projection, Bayesian identification, ...**

This approach characterizes the dependence of the first moments with respect to the parameters Q .

Another approach of GSA

Here, we exploit the structure of the model to take an alternative approach, inspired from the hierarchical orthogonal **Sobol-Hoeffding decomposition** of X :

$$X(W, Q) = \bar{X} + X_W(W) + X_Q(Q) + X_{W,Q}(W, Q), \quad \forall t \in \mathcal{T},$$

where **the functionals in the SH decomposition are mutually orthogonal**.

In fact, the decomposition is unique and given by

- $\bar{X}(t) \doteq \mathbb{E} \{X(t)\}$,
- $X_W(t, W) \doteq \mathbb{E} \{X(t)|W\} - \mathbb{E} \{X(t)\} = X_{|W}(t) - \bar{X}(t)$,
- $X_Q(t, Q) \doteq \mathbb{E} \{X(t)|Q\} - \mathbb{E} \{X(t)\} = X_{|Q}(t) - \bar{X}(t)$.

Owing to the orthogonality of the SH decomposition, we have

$$\mathbb{V} \{X\} = \mathbb{V} \{X_W\} + \mathbb{V} \{X_Q\} + \mathbb{V} \{X_{W,Q}\},$$

from which follow the definitions of the sensitivity indices

$$S_W = \frac{\mathbb{V} \{X_W\}}{\mathbb{V} \{X\}}, \quad S_Q = \frac{\mathbb{V} \{X_Q\}}{\mathbb{V} \{X\}}, \quad S_{W,Q} = \frac{\mathbb{V} \{X_{W,Q}\}}{\mathbb{V} \{X\}}.$$

Sensitivity indices

The sensitivity indices

$$S_W = \frac{\mathbb{V}\{X_W\}}{\mathbb{V}\{X\}}, \quad S_Q = \frac{\mathbb{V}\{X_Q\}}{\mathbb{V}\{X\}}, \quad S_{W,Q} = \frac{\mathbb{V}\{X_{W,Q}\}}{\mathbb{V}\{X\}},$$

then measure the fraction of the variance due to

- the **Wiener noise only**, or intrinsic randomness (S_W),
- the **parameters only**, or parametric randomness (S_Q),
- the **combined effect** of intrinsic and parametric randomness ($S_{W,Q}$).

In particular, S_W measure the part of the variance that cannot be reduced through a better knowledge of the parameters.

In addition,

$$\frac{\mathbb{V}_Q\{\mu_X(Q)\}}{\mathbb{V}\{X\}} = S_Q, \quad \text{but} \quad \frac{\mathbb{E}_Q\{\Sigma^2(Q)\}}{\mathbb{V}\{X\}} = S_W + S_{W,Q}.$$

From $\Sigma^2(Q)$, **one cannot distinguish the intrinsic and mixed randomness effects.**

Polynomial Chaos expansion

We express the dependence of X on Q as a PC expansion

$$X(t, W, Q) = \sum_{\alpha} X_{\alpha}(t, W) \psi_{\alpha}(Q),$$

where

- $\{\psi_{\alpha}\}$ is a CONS of $L^2(Q, p_Q)$,
- the expansion coefficients X_{α} are random processes.

The random processes $X_{\alpha}(t)$ are the solutions of the **coupled system of SODEs**

$$dX_{\beta}(t) = \left\langle F \left(\sum_{\alpha} X_{\alpha}(t) \psi_{\alpha}; Q \right), \psi_{\beta} \right\rangle dt + \left\langle G \left(\sum_{\alpha} X_{\alpha}(t) \psi_{\alpha}; Q \right), \psi_{\beta} \right\rangle dW,$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product in $L^2(Q, p_Q)$.

This system can be solved by MC simulation (upon truncation).

PC expansion

Assuming $\Psi_0 = 1$, it comes

$$\mathbb{E}\{X\} = \mathbb{E}\{X_0\}, \quad X_Q(Q) = \sum_{\alpha \neq 0} \mathbb{E}\{X_\alpha\} \Psi_\alpha(Q), \quad X_W(W) = X_0(W) - \mathbb{E}\{X_0\},$$

and

$$X_{W,Q}(W, Q) = \sum_{\alpha \neq 0} (X_\alpha(W) - \mathbb{E}\{X_\alpha\}) \Psi_\alpha(Q).$$

Finally, the **partial variances have for expression** :

$$\mathbb{V}\{X_Q\} = \sum_{\alpha \neq 0} \mathbb{E}\{X_\alpha\}^2, \quad \mathbb{V}\{X_W\} = \mathbb{V}\{X_0\}, \quad \mathbb{V}\{X_{W,Q}\} = \sum_{\alpha \neq 0} \mathbb{V}\{X_\alpha\}.$$

Observe :

- 1 $X_Q(Q) + \mathbb{E}\{X\} = \mu_X(Q),$
- 2 $\sum_{\alpha} \mathbb{V}\{X_\alpha\} = \sum_{\alpha} \mathbb{E}\{X_\alpha^2\} - \mathbb{E}\{X_\alpha\}^2 = \mathbb{E}_Q\{\Sigma_X^2\}.$

Linear additive system

- Consider SODE with drift and diffusion terms given by :

$$C(X, Q) = Q_1 - X \quad D(X, Q) = (\nu X + 1)Q_2$$

where Q_1 and Q_2 are independent, uniformly-distributed, random variables with mean $\mu_{1,2}$ and standard deviation $\sigma_{1,2}$.


- The orthonormal PC basis consists of tensorized Legendre polynomials.
- We use for initial condition $X(t = 0) = 0$ almost surely.

Additive Noise

Additive noise model ($\nu = 0$) with $\mu_1 = 1$, $\mu_2 = 0.1$, $\sigma_1 = \sigma_2 = 0.05$:

$$dX(Q) = (Q_1 - X(Q))dt + Q_2dW,$$


a first-order expansion suffices to exactly represent $X(Q)$.



Selected trajectories and variability ranges for $[X_k](t, W)$. The plots correspond to $k = 0, 1$ and 2 , arranged from left to right.

Multiplicative Noise – I


Multiplicative noise : $Q_1 \sim \mathcal{U}[1, 0.05]$, $Q_2 \sim \mathcal{U}[0.1, 0.05]$, $\nu = 0.2$



Sample trajectories of $[X_k]$, $0 \leq k \leq 2$. Top row : order 0, bottom row : order 1 with and decreasing order in Q_1 from left to right.

Multiplicative Noise – II



Multiplicative noise : $Q_1 \sim \mathcal{U}[1, 0.05]$, $Q_2 \sim \mathcal{U}[0.1, 0.05]$, $\nu = 0.2$



Sample trajectories of $[X_k]$, $3 \leq k \leq 14$. The total order ranges from 2 (top row) to 4 (bottom row), with and decreasing order in Q_1 from left to right.

Distribution functions


Multiplicative noise : $Q_1 \sim \mathcal{U}[1, 0.05]$, $Q_2 \sim \mathcal{U}[0.1, 0.05]$, $\nu = 0.2$





Probability density functions of the modes $[X_k]$ at $t = 10$. The modes have been centered and normalized to facilitate the comparison ; the standard Gaussian distribution is also reported for reference.

Mode correlations

Projections in the planes ($[X_k], [X_{k'}]$) of realizations of the centered and normalized solution vector \mathbf{X} at time $t = 10$, for selected indices




Conditional trajectories



Left : trajectories for samples of Q and a *fixed* realization of W


Right : trajectories for samples of W at a *fixed* value of the parameters.

SH functions




Selected trajectories of X and its SH functions.

Parametric sensitivity



Effect of Q_1 and Q_2 of the (centered) conditional mean $\mu_X(Q) = \mathbb{E}\{X | Q\} - \mathbb{E}\{X\}$ and variance $\Sigma_X^2(Q) = \mathbb{V}\{X | Q\}$ at time $t = 10$

ANOVA




Evolution of the components of the total variance. Shown are variance decompositions obtained for different values of σ_1 and σ_2

Non-linear system

Consider a system with **additive noise and non-linear drift**

$$dX = F(X)dt + \delta dW = -\gamma(X - a)(X - b)(X - c)dt + \delta dW$$

where $\delta > 0$ is an additional parameter controlling the noise level, and as before W is a Wiener process. Again the IC is $X_0 = X(t = 0)$.



Sample trajectories with $a = 10$, $b = 20$, $c = 30$, $\gamma = 0.01$, and $\delta = 1$. In all cases, the initial condition coincides with $x^0 = b$.



Parametric uncertainty

- Consider an uncertain initial condition, $Q_1 \sim \mathcal{R}[17.5, 22.5]$, and forcing amplitude, $Q_2 \sim \mathcal{R}[0.5, 1.5]$.

$$dX = F(X)dt + Q_1 dW \quad X_0 = Q_2.$$

- Q_1 and Q_2 independent.
- The PC representation is based on an adaptive multiwavelet basis expansion, which enables us to accommodate for bifurcation(s).
- The use of a non polynomial basis complicates the sensitivity analysis, but the framework is essentially unaltered.





Sample trajectories



Left plots : sample set of realizations of W , the trajectories of X (time running up) for different initial conditions and two noise levels $Q_2 = 0.65$ (top plot) and $Q_2 = 1.35$ (bottom).


Right plots show for two realizations of W (top and bottom), the trajectories of X for a random sample set of values of Q_1 and Q_2 .

MW expansion





Left : partitions of the parametric domain and surface plots for $X(t = 6, W)$ as a function of Q .

Variance decomposition



Partial variances of $X(t)$

Effect of Noise amplitude



Left : comparison of the total variances $V\{X\}$ and total noise contributions $V_{\text{noise}} + V_{\text{mix}}$ to the variance, for two expected values of $\mathbb{E}\{\delta\} = 1$ and 2 .

Right : partial variances of the stochastic process $X(t)$ for the case $\mathbb{E}\{\delta\} = 2$.

Stochastic Simulator

Work with Omar Knio and Alvaro Moraes (KAUST)

Stochastic Systems

Stochastic systems

governed by **probabilistic** evolution rules expressed by the **master equation**

$$\frac{\partial P(\mathbf{x}, t | \mathbf{x}_0, t_0)}{\partial t} = \sum_{j=1}^{K_r} [a_j(\mathbf{x} - \boldsymbol{\nu}_j) P(\mathbf{x} - \boldsymbol{\nu}_j, t | \mathbf{x}_0, t_0) - a_j(\mathbf{x}) P(\mathbf{x}, t | \mathbf{x}_0, t_0)],$$

- $P(\mathbf{x}, t | \mathbf{x}_0, t_0)$: probability of $\mathbf{X} = \mathbf{x}$ at time t , given $\mathbf{X} = \mathbf{x}_0$ at time t_0 ,
- general form $\partial P(\mathbf{x}, t | \mathbf{x}_0, t_0) / \partial t = f(P(\cdot, t | \mathbf{x}_0, t_0))$ (**Markov process**),
- $\mathbf{x}(t)$: state of the system at time t ,
- K_r reactions channels,
- propensity functions a_j and state-change vectors $\boldsymbol{\nu}_j \in \mathbb{Z}^{M_s}$.

Examples includes **Reactive Networks** (chemistry, biology), social networks, ...

- Direct resolution of the master equation is usually not an option,
- Simulations of trajectories of $\mathbf{X}(t) \sim P(\mathbf{x}, t | \mathbf{x}_0, t_0)$, using a **stochastic simulator**.

Gillespie's Algorithm

Given $\mathbf{X}(t) = \mathbf{x}$, the probability of the next reaction to occur in the $[t, t + dt)$ is

$$a_0(\mathbf{x})dt = dt \sum_{j=1}^{K_r} a_j(\mathbf{x}).$$

The time to the next reaction, τ , follows an exponential distribution with mean $1/a_0(\mathbf{x})$.

Gillespie's Algorithm :

[Gillespie, 1970's]

- 1 Set $t = t_0$, $\mathbf{X} = \mathbf{x}_0$.
- 2 Repeat until $t > T$
 - Draw $\tau \sim \exp a_0(\mathbf{X})$
 - Pick randomly $k \in \{1 \dots K_r\}$ with relative probability $p_k(a_k)$
 - update $t \leftarrow t + \tau$, $\mathbf{X} \leftarrow \mathbf{X} + \nu_k$
- 3 Return $X(T) \sim P(\mathbf{x}, t | \mathbf{x}_0, t_0)$.


From a sample set of trajectory, estimate expectation of functionals $\mathbb{E}\{g(X)\}$.

Sobol Analysis of the variance

For a given output $\mathbf{X}(t)$ of a stochastic simulator, and a functional g , we would like :

**assess the contributions of different channels (or group of channels)
on the variability of $g(\mathbf{X})$**

That is : **which channel(s)** is (are) responsible for most of the variance in $g(\mathbf{X})$?



This is **not to be confused with parametric sensitivity analyses** where one wants to estimate the sensitivity of $\mathbb{E}\{g(\mathbf{X})\}$ with respect to some parameters \mathbf{q} in the definition of the dynamics (*e.g.* propensity functions).

Sobol Analysis of the variance

- $\mathbf{N}(\omega) = (N_1, \dots, N_D)$ a set of D independent random inputs N_j ,
- $F(\mathbf{N})$ a (second-order) random functional in \mathbf{N} ,

$F(\mathbf{N})$ has a **unique orthogonal decomposition** [Sobol, 2002 ; Homma & Saltelli, 1996]

$$F(\mathbf{N}) = \sum_{\mathbf{u} \in \mathcal{D}} F_{\mathbf{u}}(\mathbf{N}_{\mathbf{u}}),$$

where \mathcal{D} is the power set of $\{1, \dots, D\}$ and $\mathbf{N}_{\mathbf{u}} = (N_{u_1}, \dots, N_{u_{|\mathbf{u}|}})$. The orthogonality condition reads

$$\mathbb{E} \{F_{\mathbf{u}} F_{\mathbf{s}}\} = \int_{\Omega} F_{\mathbf{u}}(\mathbf{N}_{\mathbf{u}}(\omega)) F_{\mathbf{s}}(\mathbf{N}_{\mathbf{s}}(\omega)) d\mu(\omega) = 0,$$

so

$$\mathbb{V} \{F\} = \sum_{\mathbf{u} \in \mathcal{D} \setminus \emptyset} \mathbb{V} \{F_{\mathbf{u}}\},$$

where $\mathbb{V} \{F_{\mathbf{u}}\}$ are called **partial variances**.

Sobol Analysis of the variance (cont...)

From the variance decomposition,

$$\mathbb{V}\{F\} = \sum_{\mathbf{u} \in \mathcal{D} \setminus \emptyset} \mathbb{V}\{F_{\mathbf{u}}\},$$

- **First order sensitivity indices** $S_{\mathbf{u}}$: fraction of the variance caused by the random inputs $\mathbf{N}_{\mathbf{u}}$ **only**

$$\mathbb{V}\{F\} S_{\mathbf{u}} = \sum_{\mathbf{s} \subseteq \mathbf{u}}^{\mathbf{s} \neq \emptyset} \mathbb{V}\{F_{\mathbf{s}}\}$$

- **Total order sensitivity indices** $T_{\mathbf{u}}$: fraction of the variance caused by the random inputs $\mathbf{N}_{\mathbf{u}}$ **and interaction**

$$\mathbb{V}\{F\} T_{\mathbf{u}} = \sum_{\mathbf{s} \in \mathcal{D}}^{(\mathbf{s} \cap \mathbf{u}) \neq \emptyset} \mathbb{V}\{F_{\mathbf{s}}\}$$

The partial variances $\mathbb{V}\{F_{\mathbf{u}}\}$ can be expressed as **conditional variances** : [Homma & Saltelli, 1996]

$$\mathbb{V}\{F_{\mathbf{u}}\} = \mathbb{V}\{\mathbb{E}\{F \mid \mathbf{N}_{\mathbf{u}}\}\} - \sum_{\substack{\mathbf{s} \in \mathcal{D} \setminus \emptyset \\ \mathbf{s} \subsetneq \mathbf{u}}} \mathbb{V}\{F_{\mathbf{s}}\},$$

or

$$\mathbb{V}\{F\} S_{\mathbf{u}} = \mathbb{V}\{\mathbb{E}\{F \mid \mathbf{N}_{\mathbf{u}}\}\}, \quad \mathbb{V}\{F\} T_{\mathbf{u}} = \mathbb{V}\{F\} - \mathbb{V}\{\mathbb{E}\{F \mid \mathbf{N}_{\mathbf{u}^c}\}\} = \mathbb{V}\{F\} (1 - S_{\mathbf{u}^c}),$$

where $\mathbf{u}^c = \{1, \dots, D\} \setminus \mathbf{u}$.

Decomposition of the Variance = Estimation of conditional variances

Monte-Carlo estimation of the sensitivity indices

Consider **two independent sample sets** \mathcal{N}^I and \mathcal{N}^{II} of M realizations of \mathbf{N} .

The conditional variance $\mathbb{V}\{\mathbb{E}\{F \mid \mathbf{N}_{\mathbf{u}}\}\}$ can be estimated as

[Sobol, 2001]

$$\mathbb{V}\{\mathbb{E}\{F \mid \mathbf{N}_{\mathbf{u}}\}\} + \mathbb{E}\{F\}^2 = \lim_{M \rightarrow \infty} \frac{1}{M} \sum_{i=1}^M F(\mathbf{N}_{\mathbf{u}}^{I,(i)}, \mathbf{N}_{\mathbf{u}\sim}^{I,(i)}) F(\mathbf{N}_{\mathbf{u}}^{II,(i)}, \mathbf{N}_{\mathbf{u}\sim}^{II,(i)}),$$

such that

$$\widehat{S}_{\mathbf{u}} = \frac{\frac{1}{M} \sum_{i=1}^M F(\mathbf{N}^{I,(i)}) F(\mathbf{N}_{\mathbf{u}}^{I,(i)}, \mathbf{N}_{\mathbf{u}\sim}^{I,(i)}) - \widehat{\mathbb{E}\{F\}}^2}{\widehat{\mathbb{V}\{F\}}},$$

and

$$\widehat{T}_{\mathbf{u}} = 1 - \frac{\frac{1}{M} \sum_{i=1}^M F(\mathbf{N}^{II,(i)}) F(\mathbf{N}_{\mathbf{u}}^{II,(i)}, \mathbf{N}_{\mathbf{u}\sim}^{II,(i)}) - \widehat{\mathbb{E}\{F\}}^2}{\widehat{\mathbb{V}\{F\}}}$$

where $\widehat{\mathbb{E}\{F\}}$ and $\widehat{\mathbb{V}\{F\}}$ are the classical MC estimators for the mean and variance.

The computational complexity scales linearly with the number of indices to be computed

Application to Stochastic Simulators

To assess the **respective impacts of different reaction channels** through Sobol's decomposition of $\mathbb{V}\{g(\mathbf{X})\}$, when \mathbf{X} is the output of a stochastic simulator, we need to **condition \mathbf{X} on the channels dynamics** :

What is a particular realization of a channel dynamics ?

Gillespie's algorithm is not suited, and we have to recast the stochastic algorithm in terms of

independent processes associated to each channel.

Next Reaction Formulation.

[Ethier & Kurtz, 2005, Gibson & Bruck, 2000]

$$\mathbf{X}(t) = \mathbf{X}(t_0) + \sum_{j=1}^{K_r} \nu_j N_j(t_j),$$

where the $N_j(t)$ are independent standard (unit rate) Poisson processes, and the scaled times t_j are given by

$$t_j = \int_{t_0}^t a_j(\mathbf{X}(\tau)) d\tau, \quad j = 1, \dots, K_r.$$

Then, $g(\mathbf{X})$ can be seen as

$$g(\mathbf{X}) = F(N_1, \dots, N_{K_r}).$$

Application to Stochastic Simulators (cont.)

The random functional $g(\mathbf{X}) = F(N_1, \dots, N_{K_r})$ can then be decomposed *à la* Sobol.

A particular **realization of a channel dynamic is identified with a realization of the underlying standard Poisson processes.**

For instance, the conditional variance writes

$$\mathbb{E} \{ g(\mathbf{X}) \mid \mathbf{N}_{\mathbf{u}} = \mathbf{n}_{\mathbf{u}} \} = \mathbb{E} \left\{ g \left(\mathbf{X}(t_0) + \sum_{j \in \mathbf{u}} \nu_j n_j(t_j) + \sum_{j \in \mathbf{u}_{\sim}} \nu_j N_j(t_j) \right) \right\},$$

with $t_j = \int_{t_0}^t a_j(\mathbf{X}(\tau)) d\tau$.


Note that

- in general, for fixed $\mathbf{n}_{\mathbf{u}}(t)$ the **physical firing times t_j remain random** and depend on $\mathbf{N}_{\mathbf{u}_{\sim}}$, even for $j \in \mathbf{u}$
- in practice, the standard Poisson processes N_j are entirely specified by their random seeds and pseudo-random number generator :

the Poisson processes don't have to be stored but are computed on the fly

The birth-death (BD) process

Single species S ($M_S = 1$) and $K_r = 2$ reaction channels :



with propensity functions

$$a_1(x) = b, \quad a_2(x) = d \times x.$$

We set $b = 200$, $d = 1$, and use $M = 1,000,000$ Monte Carlo samples to compute the estimates.







FIGURE: Left : Selected trajectories of $X(t)$ generated using Next Reaction Algorithm. Right : histogram of $X(t = 8)$.


B-D process. Variance decomposition of $g(X) = X(t)$ 

Left : scaled first-order and total sensitivity indices (scaled by the variance) of the birth-death model and $t \in [0, 8]$. Right : long-time evolution of the first-order sensitivity indices, and of the mixed interaction term.

- Variance in X is predominantly caused by the birth channel stochasticity for early time $t < 1$
- For $1 \leq t \leq 4$, the variability induced by R_d only continues to grow with the population size (first order reaction), while mixed effects develops
- Eventually, effect of R_b stabilize (zero-order reaction, rate independent of X) while effect of R_d only slowly decays to benefit the mixed term (stochasticity of N_b affects more and more the death process).

Schlögl model

System with $K_r = 4$ reaction channels :




with B_1 and B_2 in large excess and constant population over time, $X_{B_1} = X_{B_2}/2 = 10^5$ and a **single evolving species S** with $M_S = 1$. The propensity functions are given by

$$a_1(x) = \frac{c_1}{2} X_{B_1} x(x-1), \quad a_2(x) = \frac{c_2}{6} x(x-1)(x-2), \quad a_3(x) = c_3 X_{B_2}, \quad a_4(x) = c_4 x.$$


We set $c_1 = 3 \times 10^{-7}$, $c_2 = 10^{-4}$, $c_3 = 10^{-3}$, $c_4 = 3.5$ and deterministic initial condition $X(t=0) = 250$.

Results in a bi-modal dynamic





Left : selected trajectories of $X(t)$ showing the bifurcation in the stochastic dynamics. Right : histogram of $X(t=8)$.

Schlögl model - Variance decomposition of $g(X) = X(t)$



First and total order partial variances.




Higher order partial variances.

Reaction channels R_1 and R_4 are the dominant sources of variance
 Dynamic essentially additive up to $t \sim 2$

Schlögl model - Variance decomposition of $g(X) = X(t)$

Analysis of the partial variance revealed that R_1 and R_4 are the main sources of stochasticity.


It suggests a **dominant role in selecting the bifurcation branch**, as illustrated below



Trajectories of $X(t)$ **conditioned** on (a) $N_1(\omega) = n_1$ and $N_4(\omega) = n_4$, and (b) $N_2(\omega) = n_2$ and $N_3(\omega) = n_3$. Each sub-plot shows 10 conditionally random trajectories for fixed realizations n_1 and n_4 in (a), and n_2 and n_3 in (b).


Michaelis-Menten system

$M_s = 4$ species and $K_r = 3$ reaction channels :




with $a_1(\mathbf{x}) = c_1 x_1 x_2$, $a_2(\mathbf{x}) = c_2 x_3$, and $a_3(\mathbf{x}) = c_3 x_3$,

We set $c_1 = 0.0017$, $c_2 = 10^{-3}$ and $c_3 = 0.125$, and initial conditions
 $X_1(t=0) = 300$, $X_2(t=0) = 120$ and $X_3(t=0) = X_4(t=0) = 0$



Michaelis-Menten system - Variance decomposition of $g(\mathbf{X}) = X_1(t)$

Note : $X_2 + X_3 = \text{const}$, the sensitivity indices for S_2 and S_3 are equal




Michaelis-Menten model : First-order and total sensitivity indices $S_{\{j\}}$ and $T_{\{j\}}$ for $j = 1, \dots, 4$.
Plots are generated for (a) X_1 , (b) X_2 and (c) X_4

- Relative importance of R_1 and R_3 changes in time for S_1 and S_2
- Stochastic dynamic of S_4 is essentially additive and dominated by R_3
- Channel R_2 induces nearly no variance in $X(t)$: **here** the dissociation reaction R_2 can be simply disregarded without affecting significantly the dynamics.

Michaelis-Menten system - Variance decomposition of $g(\mathbf{X}) = X_i(t)$

On the contrary, increasing c_2 by an order of magnitude, the effect of R_2 on the variances becomes apparent :



Conclusion and Future Work

We have proposed

- Hybrid Galerkin / MC approach for noisy systems, can be extended to Non-Intrusive methods
- Parametric dependence assumes **smoothness of the trajectories**
- Development of **methods and algorithms** that enable variance decomposition for stochastic simulators output
- Identification of the channels dynamics with their underlying standard Poisson processes
- Assessment of the **relative importance of different reaction channels** on the variance

On-going and future works

- Application to more complex functional $g(\mathbf{X})$: exit-time, path integrals, ... (smoothness ?)
- Account for parametric uncertainty in the definition of the propensity functions
- Improve stochastic simulators for computational complexity reduction, *e.g.* Tau-Leaping method and variance reduction methods.

[OLM & OK, J. Rel. Eng. Sys. Safety, 2015], [OLM, OK and A. Moraes, J. Chem. Phys., 2015 in revision]

Acknowledgement : This work was supported by the US Department of Energy (DOE), Office of Science, Office of Advanced Scientific Computing Research, under Award Number DE-SC0008789. Support from the Research Center on Uncertainty Quantification of the King Abdullah University of Science and Technology is also acknowledged.

Algorithm

ALGORITHM 3. Computation of the first and total-order sensitivity indices $S_{(j)}$ and $T_{(j)}$ of $g(\mathbf{X}(T))$.

Procedure Compute_SI($M, \mathbf{X}_0, T, \{\nu_j\}, \{a_j\}, g$)

Require: Sample set dimension M , initial condition \mathbf{X}_0 , final time T , state-change vectors $\{\nu_j\}$, propensity functions $\{a_j\}$ and functional g

```

1:  $\mu \leftarrow 0, \sigma^2 \leftarrow 0$             $\triangleright$  Init. Mean and Variance
2: for  $j = 1$  to  $K_r$  do
3:    $S(j) \leftarrow 0, T(j) \leftarrow 0$     $\triangleright$  Init. first and total-order SIs
4: end for
5: for  $m = 1$  to  $m = M$  do
6:   Draw two independent set of seeds  $\mathbf{s}^I$  and  $\mathbf{s}^{II}$ 
7:    $\mathbf{X} \leftarrow \text{NRA}(\mathbf{X}_0, T, \{\nu_j\}, \{a_j\}, \text{RG}_1(\mathbf{s}^I), \dots, \text{RG}_{K_r}(\mathbf{s}^{II}_{K_r}))$ 
8:    $\mu \leftarrow \mu + g(\mathbf{X}), \sigma^2 \leftarrow \sigma^2 + g(\mathbf{X})^2$     $\triangleright$  Acc. mean and variance
9:   for  $j = 1$  to  $K_r$  do
10:     $\mathbf{X}_S \leftarrow \text{NRA}(\mathbf{X}_0, T, \{\nu_j\}, \{a_j\}, \text{RG}_1(\mathbf{s}^{II}_1), \dots,$ 
11:       $\dots, \text{RG}_j(\mathbf{s}^{II}_j), \dots, \text{RG}_{K_r}(\mathbf{s}^{II}_{K_r}))$ 
12:     $\mathbf{X}_T \leftarrow \text{NRA}(\mathbf{X}_0, T, \{\nu_j\}, \{a_j\}, \text{RG}_1(\mathbf{s}^I_1), \dots,$ 
13:       $\dots, \text{RG}_j(\mathbf{s}^I_j), \dots, \text{RG}_{K_r}(\mathbf{s}^I_{K_r}))$ 
14:     $S(j) \leftarrow S(j) + g(\mathbf{X}) \times g(\mathbf{X}_S)$     $\triangleright$  Acc. 1-st order
15:     $T(j) \leftarrow T(j) + g(\mathbf{X}) \times g(\mathbf{X}_T)$     $\triangleright$  Acc. total order
16:  end for    $\triangleright$  Next channel
17: end for    $\triangleright$  Next sample
18:  $\mu \leftarrow \mu/M, \sigma^2 \leftarrow \sigma^2/(M-1) - \mu^2$ 
19: for  $j = 1$  to  $K_r$  do
20:    $S(j) \leftarrow \frac{S(j)}{(M-1)\sigma^2} - \frac{\mu^2}{\sigma^2}$     $\triangleright$  Estim. 1-st order
21:    $T(j) \leftarrow 1 - \frac{T(j)}{(M-1)\sigma^2} + \frac{\mu^2}{\sigma^2}$     $\triangleright$  Estim. total order
22: end for
23: Return  $S(j)$  and  $T(j), j = 1, \dots, K_r$     $\triangleright$  First and total-order sensitivity indices  $S_{(j)}$  and  $T_{(j)}$  of  $g(\mathbf{X}(T))$ 

```

- Procedure NRA implement the Next Reaction Algorithm
- Poisson processes defined by two independent sets of seeds and RNG
- Obvious parallelization

ALGORITHM 2. Next Reaction Algorithm.

Procedure NRA($\mathbf{X}_0, T, \{\nu_j\}, \{a_j\}, \text{RG}_1, \dots, \text{RG}_{K_r}$)

Require: Initial condition \mathbf{X}_0 , final time T , state-change vectors $\{\nu_j\}$, propensity functions $\{a_j\}$, and seeded pseudo-random number generators $\text{RG}_{j=1, \dots, K_r}$.

```

1: for  $j = 1, \dots, K_r$  do
2:   Draw  $r_j$  from  $\text{RG}_j$ 
3:    $\tau_j \leftarrow 0, \tau_j^+ \leftarrow -\log r_j$     $\triangleright$  set next reaction times
4: end for
5:  $t \leftarrow 0, \mathbf{X} \leftarrow \mathbf{X}_0$ 
6: loop
7:   for  $j = 1, \dots, K_r$  do
8:     Evaluate  $a_j(\mathbf{X})$  and  $dt_j = \frac{\tau_j^+ - \tau_j}{a_j}$ 
9:   end for
10:  Set  $l = \arg \min_j dt_j$     $\triangleright$  pick next reaction
11:  if  $t + dt_l > T$  then
12:    break    $\triangleright$  Final time reached
13:  else
14:     $t \leftarrow t + dt_l$     $\triangleright$  update time
15:     $\mathbf{X} \leftarrow \mathbf{X} + \nu_l$     $\triangleright$  update the state vector
16:    for  $j = 1, \dots, K_r$  do
17:       $\tau_j \leftarrow \tau_j + a_j dt_l$     $\triangleright$  update unscaled times
18:    end for
19:    Get  $r_l$  from  $\text{RG}_l$ 
20:     $\tau_l^+ \leftarrow \tau_l^+ - \log r_l$     $\triangleright$  next reaction time
21:  end if
22: end loop
23: Return  $\mathbf{X}$     $\triangleright$  State  $\mathbf{X}(T)$ 

```